



Fracture Behavior with Cheese

Learning Objectives	<ol style="list-style-type: none"> 1) Students will apply their knowledge of fracture to predict how different cheeses will fail. 2) Students will discuss how geometry plays a role in fracture behavior.
Time	~60 minutes
Topics	<ol style="list-style-type: none"> 1) Mechanical testing to characterize fracture properties 2) Stress concentrators 3) Simulations of stress (computational materials science)

Before the Lab:

Supplies to buy every time you run the activity:

- 1) Two different types of cheese. Choose cheese with different types of fracture.

Cheese Type	Fracture Type
Mozzarella	Ductile
Feta	Intergranular
Parmesan	Cleavage
Colby Jack	Ductile
Muenster	Ductile
Extra Sharp Cheddar	Cleavage

Supplies to buy as needed:

- 1) Paper plates for the cheese
- 2) Gloves for those who are lactose intolerant
- 3) A knife and toothpick to make the notches in the cheese

Prior Knowledge

Recommended for Instructor:

- 1) Familiarity with basic mechanical properties
 - Stress vs. strain graphs for brittle vs. ductile materials
 - Three-point bend testing
- 2) Basic understanding of material fracture
 - Understand the difference between different types of fracture mechanisms

Prior Knowledge

Recommended for Students:

- 1) Types of bonding
 - Covalent, non-covalent, ionic



Fracture Behavior with Cheese

You Will Be Able To:

Explain why geometry of a material is important to understanding how it fractures

Files Needed for Simulation:

Preload these into an easily identifiable file prior to the exercise if possible

Blunt_crack.png	Narrow_crack_script.py	No_crack.png
Blunt_crack_script.py	Center_crack.png	No_crack_script.py
Narrow_crack.png	Center_crack_script.py	

Simulations of Stress in Cheese

When a material breaks, there are engineers who will figure out why it broke. Instead of breaking a bunch of objects to see why they failed, engineers use different kinds of computer programs to digitally model how a material would fail. You will use one of these programs, called OOF2, to understand why the geometry of a material can concentrate physical stress in a particular location in the material. Each group will be assigned to simulate a different initial fracture geometry (due to time constraints), which you will present at the end. What you will see once you have finished the simulation is a visual representation of where stress builds up in the cheese as it is initially strained.

Notes to the instructor: Nanohub, and OOF2 especially, can be kind of finicky.

Try to use a browser with javascript support (Safari and Firefox are usually better than Chrome), and if the display cuts out, try refreshing the page before assuming the worst. Sometimes it really will just break for no adequately explained reason and the simulation will have to be rerun (this is the case if there's a connection error of some kind, in our experience.)

If refreshing the page doesn't work you can try going to the student's "dashboard" and launching from currently-running sessions.

One other thing to note is that some of these simulations could take 5-10 minutes to fully set up and solve, especially ones with more complex meshes.


If the particular crack file will not load correctly just have them run a different crack.

Fracture Behavior with Cheese


Example Solution Images:

These won't look identical from group to group because there are random steps in the OOF2 setup, but they should be a solid guide for what to look at. Note that if they don't uncheck their original image file, their images may not appear to have as much contrast as these do. Also, generally these can be most easily created by just screenshotting the Nanohub images. Note also that the scales on the color bars are different, which may be relevant.


Blunt Crack



Center Crack




Narrow Crack



Fracture Behavior with Cheese

Example Solution Images (cont.)

No Crack



- **No crack:** the color variations are just noise as the whole field is subjected to a uniform stress
- **Blunt crack:** smallest range of tensile to compressive stresses, with the most gradual changes between contours – likely to result in more ductile fracture at higher loads
- **Center crack:** much larger range of stress and much more localized
- **Narrow crack:** highly localized fields at the crack tip, the crack tip radius and the loading direction the most relevant parameters

Discussion Points:

What you are looking at is a scaled color map of the stress intensities at various locations around an initial fracture shape (or no shape, if you are simulating without an initial crack)

1. Using this picture, predict how you think a soft cheese like mozzarella would fracture compared to a harder cheese like parmesan for different types of crack geometries?

See summary above for description and rationale of stress concentration fields. When a crack propagates:

- **no crack** will nucleate and fail at random points along the specimen;
- **blunt crack** is likely to have a zig-zag crack path (ductile tearing, especially in the “softer” cheese);
- **center crack** should propagate on the tensile side of the bending specimen first, but if loaded in pure tension (as demonstrated in this simulation) both sides should fail simultaneously in much the same way that the sharp crack would although at different loads
- **narrow crack** should propagate in a zig-zag pattern for softer cheese and in a straight path perpendicular to the crack tip for harder cheese

2. What kind of fracture behavior do you think you have modeled? Explain.

As noted above, this depends not only on the crack geometry but also on the stiffness


3. Compare your simulation results to the results of the other groups. From the simulations, how does initial geometry play a role in crack propagation?

Guide the campers towards the conclusions noted above

Fracture Behavior with Cheese

Lab Set-up:

- 1) Break students up into pairs. Half the groups will receive Extra Sharp Cheddar Cheese and half the groups will receive Muenster cheese. Arrange the groups so that groups with different cheeses are sitting close to each other to encourage sharing their results after the experiment.
- 2) Before the students arrive, make sure each group has:
 - One plate with five cheese samples. Each sample should be approximately 2 inches x 1 inch.
 - Two cheese samples with sharp notch
 - Two cheese samples with blunt notch
 - One cheese sample with no notch
 - Gloves (if requested by the participant)
- 3) Pre-notch all of the cheese samples (see three-point bend test figure above). The sharp notch is made using a knife to make a small slice. The blunt notch is made using a knife to cut a small wedge and then a toothpick to smooth out the edges.




Pre-lab Questions:

The lecture before this lab should explain the basics of fracture and the types of fracture mechanisms.


- 1) The cheese fracture activity fits into the structure and properties categories of the tetrahedron. Please explain why cheese fracture fits in these categories.

Cheese fits into the structure and properties categories because the type of bonding in the cheese affects the way it fractures, which is a mechanical property.




- 2) Match the type of fracture to its fracture surface


Intergranular



Ductile Fracture



Cleavage





Fracture Behavior with Cheese

Running the Lab:

- 1) Have the students sit in pairs at their stations. Explain they will be handling cheese. Offer gloves to those who are lactose intolerant or would prefer not to handle cheese with their bare hands.
- 2) They should have already answered the pre-lab questions. Ask the students if they had any unresolved questions about the pre-lab questions.
- 3) Instruct the students to read all of the directions.
- 4) Do not give them the instructions verbally. The goal is to get the students to read everything and think about why they are doing what they are doing. At this point let them jump in but encourage them to raise their hands if they have questions.

Expected Results from Three-point Bend Tests:

We provide students with more than one of the same geometry because there are often defects that impact the way the specimens fail. By inspecting more than one specimen, students can also explore the real-life variability of failure analysis. The chart below is completed to provide the instructors with some insight about both types of cheese in each geometry; however, students would only be working with one type at a time. The examples below do not include any descriptions of intergranular fracture. We have found that these cheeses (e.g. feta or Colby jack) can be a little frustrating for students. Instead, we suggest that you provide a few examples to demonstrate intergranular fracture.

Sample	Crack nucleation	Crack propagation (Straight or zig-zag?)	Fracture surface (Ductile Fracture, Cleavage or Intergranular?)
Sharp notch #1 (for Extra Sharp Cheddar)	<i>The crack should start directly from the sharp notch.</i>	<i>Most cracks will propagate directly across the specimen (i.e. straight)</i>	<i>Extra sharp cheddar should demonstrate cleavage-like behavior with signature "river-pattern" markings</i>
Sharp notch #2 (for Muenster)	<i>The crack should start directly from the sharp notch.</i>	<i>Most cracks will propagate at an angle to the original notch (~45°). Often this results in a more jagged or zig-zag propagation path.</i>	<i>Muenster cheese typically displays ductile fracture behavior. The zig-zag crack path and void formation ahead of the crack are signatures of this.</i>
Blunt notch #1 (for Extra Sharp Cheddar)	<i>The crack is still expected to start from the notch.</i>	<i>While there may be more variation than found with the sharp cracks, most will still propagate straight across the specimen.</i>	<i>The fracture surface should still have signatures of brittle cleavage fracture.</i>
Blunt notch #2 (for Muenster)	<i>The crack may start at the notch, but is just as likely to start somewhere else.</i>	<i>This crack path is still likely to be zig-zaggy, but will also be the most unpredictable.</i>	<i>Still ductile fracture.</i>
No notch	<i>For either type of cheese, without a notch, the crack can start anywhere.</i>	<i>Without a pre-notch, the crack is much more difficult to predict and the students will likely have to apply more force to initiate one.</i>	<i>This will depend on the type of cheese, but the signatures of fracture should be the same as observed with the pre-notched samples.</i>

Discussion Points:

- 1) What does your fracture surface look like for each cheese? For each type of notch?
- 2) Did all of your samples fail the same way? Can you rationalize why or why not?